

Product datasheet for TA356537

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Transferrin Receptor 2 (TFR2) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications:IHC, WBReactivity:HumanHost:Rabbit

Clonality: Polyclonal

Immunogen: The immunogen is a synthetic peptide directed towards the N terminal region of human TFR2

Specificity: Expected reactivity: Cow, Dog, Guinea Pig, Horse, Human, Mouse, Rabbit, Rat

Homology: Cow: 100%; Dog: 100%; Guinea Pig: 100%; Horse: 100%; Human: 100%; Mouse:

100%; Rabbit: 100%; Rat: 100%

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

Concentration: lot specific

Purification: Affinity Purified
Conjugation: Unconjugated

Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small

aliquots to prevent freeze-thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 89kDa

Gene Name: transferrin receptor 2

Database Link: NP 003218

Entrez Gene 7036 Human

Q9UP52






Background:

TFR2,a member of the transferrin receptor-like family,is a single-pass type II membrane protein with a protease associated (PA) domain, an M28 peptidase domain and a transferrin receptor-like dimerization domain. This protein mediates cellular uptake of transferrin-bound iron and mutations in this gene have been associated with hereditary hemochromatosis type III. This gene is a member of the transferrin receptor-like family and encodes a single-pass type II membrane protein with a protease associated (PA) domain, an M28 peptidase domain and a transferrin receptor-like dimerization domain. This protein mediates cellular uptake of transferrin-bound iron and mutations in this gene have been associated with hereditary hemochromatosis type III. Alternatively spliced variants which encode different protein isoforms have been described; however, not all variants have been fully characterized. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Entrez Gene record to access additional publications.

Synonyms: HFE3; MGC126368; TFRC2

Protein Families: Druggable Genome, Protease, Transmembrane

Product images:




WB Suggested Anti-TFR2 Antibody Titration: 0.2-1

ug/ml

ELISA Titer: 1:62500


Positive Control: Human Muscle





Host: Rabbit Target Name: TFR2 Positive Control: ~25ug HepG2 Cell Lysate (HG) Negative Control: ~25ug 293T Cell Lysate (2T) Antibody Concentration: 5ug/ml Host: Rabbit Target: TFR2

Positive control (+): HepG2 Cell Lysate (HG) Negative control (-): 293T Cell Lysate (2T) Antibody concentration: 5ug/ml



Human Skin