

# **Product datasheet for TA357611**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Metabotropic Glutamate Receptor 2 (GRM2) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WB

Reactivity: Human Host: Rabbit

Clonality: Polyclonal

**Immunogen:** The immunogen is a synthetic peptide directed towards the C-terminal region of Human

GRM2

Specificity: Expected reactivity: Cow, Dog, Guinea Pig, Horse, Human, Mouse, Rabbit, Rat

**Homology**: Cow: 100%; Dog: 100%; Guinea Pig: 100%; Horse: 100%; Human: 100%; Mouse:

93%; Rabbit: 100%; Rat: 100%

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

**Concentration:** lot specific

Purification: Affinity purified Conjugation: Unconjugated

Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small

aliquots to prevent freeze-thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 95 kDa

**Gene Name:** glutamate metabotropic receptor 2

**Database Link:** XP 006713184.1

Entrez Gene 2912 Human

Q14416






#### Background:

L-glutamate is the major excitatory neurotransmitter in the central nervous system and activates both ionotropic and metabotropic glutamate receptors. Glutamatergic neurotransmission is involved in most aspects of normal brain function and can be perturbed in many neuropathologic conditions. The metabotropic glutamate receptors are a family of G protein-coupled receptors, that have been divided into 3 groups on the basis of sequence homology, putative signal transduction mechanisms, and pharmacologic properties. Group I includes GRM1 and GRM5 and these receptors have been shown to activate phospholipase C. Group II includes GRM2 and GRM3 while Group III includes GRM4, GRM6, GRM7 and GRM8. Group II and III receptors are linked to the inhibition of the cyclic AMP cascade but differ in their agonist selectivities. Two transcript variants encoding different isoforms have been found for this gene.

Synonyms: GLUR2; GPRC1B; mGlu2; MGLUR2

# **Product images:**



WB Suggested Anti-GRM2 antibody Titration: 1

ug/mL

Sample Type: Human Uterus Tumor