

# **Product datasheet for TA358004**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Silencer of Death Domain (BAG4) Rabbit Polyclonal Antibody

#### **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Reactivity: Human Host: Rabbit

**Clonality:** Polyclonal

**Specificity: Expected reactivity**: Cow, Dog, Guinea Pig, Horse, Human, Mouse, Rabbit, Rat, Zebrafish

**Homology**: Cow: 100%; Dog: 100%; Guinea Pig: 93%; Horse: 100%; Human: 100%; Mouse:

100%; Rabbit: 93%; Rat: 100%; Zebrafish: 92%

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

**Concentration:** lot specific

Purification: Affinity Purified Conjugation: Unconjugated

Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small

aliquots to prevent freeze-thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 50kDa

**Gene Name:** BCL2 associated athanogene 4

Database Link: NP 004865

Entrez Gene 9530 Human

095429






Background:

The protein encoded by this gene is a member of the BAG1-related protein family. BAG1 is an anti-apoptotic protein that functions through interactions with a variety of cell apoptosis and growth related proteins including BCL-2, Raf-protein kinase, steroid hormone receptors, growth factor receptors and members of the heat shock protein 70 kDa family. This protein contains a BAG domain near the C-terminus, which could bind and inhibit the chaperone activity of Hsc70/Hsp70. This protein was found to be associated with the death domain of tumor necrosis factor receptor type 1 (TNF-R1) and death receptor-3 (DR3), and thereby negatively regulates downstream cell death signaling. The regulatory role of this protein in cell death was demonstrated in epithelial cells which undergo apoptosis while integrin mediated matrix contacts are lost. Alternatively spliced transcript variants encoding distinct isoforms have been identified.

Synonyms: BAG-4; SODD

**Protein Families:** Druggable Genome

## **Product images:**



WB Suggested Anti-BAG4 Antibody Titration: 1.0 ug/ml

Positive Control: RPMI-8226 Whole Cell