

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA358471

RPL22 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	Expected reactivity : Cow, Dog, Guinea Pig, Horse, Human, Mouse, Rabbit, Rat, Zebrafish Homology : Cow: 100%; Dog: 100%; Guinea Pig: 100%; Horse: 100%; Human: 100%; Mouse: 100%; Rabbit: 100%; Rat: 100%; Zebrafish: 100%
Formulation:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Concentration:	lot specific
Purification:	Affinity Purified
Conjugation:	Unconjugated
Storage:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	14kDa
Gene Name:	ribosomal protein L22
Database Link:	<u>NP_000974</u> <u>Entrez Gene 6146 Human</u> <u>P35268</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE RPL22 Rabbit Polyclonal Antibody – TA358471


Background:Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and
a large 60S subunit. Together these subunits are composed of 4 RNA species and
approximately 80 structurally distinct proteins. This gene encodes a cytoplasmic ribosomal
protein that is a component of the 60S subunit. The protein belongs to the L22E family of
ribosomal proteins. Its initiating methionine residue is post-translationally removed. The
protein can bind specifically to Epstein-Barr virus-encoded RNAs (EBERs) 1 and 2. The mouse
protein has been shown to be capable of binding to heparin. Transcript variants utilizing
alternative polyA signals exist. As is typical for genes encoding ribosomal proteins, there are
multiple processed pseudogenes of this gene dispersed through the genome. It was
previously thought that this gene mapped to 3q26 and that it was fused to the acute myeloid
leukemia 1 (AML1) gene located at 21q22 in some therapy-related myelodysplastic syndrome
patients with 3;21 translocations; however, these fusions actually involve a ribosomal protein
L22 pseudogene located at 3q26, and this gene actually maps to 1p36.3-p36.2.

Synonyms: EAP; HBP15; HBP15/L22

Ribosome

Protein Pathways:

Product images:



WB Suggested Anti-RPL22 Antibody Titration: 1.0 ug/ml Positive Control: Placenta

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US