

Product datasheet for **TA358618**

CDw75 (ST6GAL1) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Specificity:	Expected reactivity: Cow, Dog, Goat, Guinea Pig, Horse, Human, Rabbit, Rat, Sheep Homology: Cow: 92%; Dog: 100%; Goat: 92%; Guinea Pig: 100%; Horse: 100%; Human: 100%; Rabbit: 100%; Rat: 100%; Sheep: 92%
Formulation:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. <i>Note that this product is shipped as lyophilized powder to China customers.</i>
Concentration:	lot specific
Purification:	Affinity Purified
Conjugation:	Unconjugated
Storage:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	46kDa
Gene Name:	ST6 beta-galactoside alpha-2,6-sialyltransferase 1
Database Link:	NP_003023 Entrez Gene 6480 Human P15907
Background:	This gene encodes a member of glycosyltransferase family 29. The encoded protein is a type II membrane protein that catalyzes the transfer of sialic acid from CMP-sialic acid to galactose-containing substrates. The protein, which is normally found in the Golgi but can be proteolytically processed to a soluble form, is involved in the generation of the cell-surface carbohydrate determinants and differentiation antigens HB-6, CD75, and CD76. This gene has been incorrectly referred to as CD75. Three transcript variants encoding two different isoforms have been described.




[View online »](#)

Synonyms: CMP-N-acetylneuraminate-beta-galactosamide-alpha-2; MGC48859; OTTHUMP00000210002; SIAT1; ST6GalI; ST6N

Protein Families: Secreted Protein

Protein Pathways: Metabolic pathways, N-Glycan biosynthesis

Product images:



WB Suggested Anti-ST6GAL1 Antibody
Titration: 1.0 ug/ml
Positive Control: HepG2 Whole Cell
ST6GAL1 is strongly supported by BioGPS gene expression data to be expressed in Human HepG2 cells