

## **Product datasheet for TA358623**

## **DUSP4 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type: Primary Antibodies** 

**Applications:** WB

Reactivity: Human Rabbit Host:

Clonality: Polyclonal

Specificity: Expected reactivity: Cow, Dog, Guinea Pig, Horse, Human, Mouse, Pig, Rabbit, Rat

Homology: Cow: 92%; Dog: 85%; Guinea Pig: 86%; Horse: 92%; Human: 100%; Mouse: 92%;

Pig: 92%; Rabbit: 85%; Rat: 92%

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

Concentration: lot specific

**Purification: Affinity Purified** Conjugation: Unconjugated

Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small

aliquots to prevent freeze-thaw cycles.

Shelf life: one year from despatch. Stability:

**Predicted Protein Size:** 43kDa

Gene Name: dual specificity phosphatase 4

Database Link: NP 001385

Entrez Gene 1846 Human

Q13115



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:


The protein encoded by this gene is a member of the dual specificity protein phosphatase subfamily. These phosphatases inactivate their target kinases by dephosphorylating both the phosphoserine/threonine and phosphotyrosine residues. They negatively regulate members of the mitogen-activated protein (MAP) kinase superfamily (MAPK/ERK, SAPK/JNK, p38), which are associated with cellular proliferation and differentiation. Different members of the family of dual specificity phosphatases show distinct substrate specificities for various MAP kinases, different tissue distribution and subcellular localization, and different modes of inducibility of their expression by extracellular stimuli. This gene product inactivates ERK1, ERK2 and JNK, is expressed in a variety of tissues, and is localized in the nucleus. Two alternatively spliced transcript variants, encoding distinct isoforms, have been observed for this gene. In addition, multiple polyadenylation sites have been reported.

Synonyms: HVH2; MKP-2; MKP2; TYP; VH2

**Protein Families:** Phosphatase

**Protein Pathways:** MAPK signaling pathway

## **Product images:**



WB Suggested Anti-DUSP4 Antibody

Titration: 1.0 ug/ml

Positive Control: COLO205 Whole CellDUSP4 is strongly supported by BioGPS gene expression data to be expressed in Human COLO205 cells