

Product datasheet for TA359378

BRSK2 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Reactivity: Human Host: Rabbit

Clonality: Polyclonal

Immunogen: The immunogen is a synthetic peptide directed towards the middle region of Human BRSK2

Expected reactivity: Human Specificity:

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

Concentration: lot specific

Purification: Affinity Purified Unconjugated Conjugation:

For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small Storage:

aliquots to prevent freeze-thaw cycles.

Shelf life: one year from despatch. Stability:

Predicted Protein Size: 84kDa

Gene Name: BR serine/threonine kinase 2

Database Link: Entrez Gene 9024 Human

O8IWO3-5



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

BRSK2 is a serine/threonine-protein kinase that plays a key role in polarization of neurons and axonogenesis, cell cycle progress and insulin secretion. It phosphorylates CDK16, CDC25C, MAPT/TAU, PAK1 and WEE1. Following phosphorylation and activation by STK11/LKB1, acts as a key regulator of polarization of cortical neurons, probably by mediating phosphorylation of microtubule-associated proteins such as MAPT/TAU at 'Thr-529' and 'Ser-579'. It also regulates neuron polarization by mediating phosphorylation of WEE1 at 'Ser-642' in post-mitotic neurons, leading to down-regulate WEE1 activity in polarized neurons. It plays a role in the regulation of the mitotic cell cycle progress and the onset of mitosis. It plays a role in the regulation of insulin secretion in response to elevated glucose levels, probably via phosphorylation of CDK16 and PAK1. While BRSK2 is phosphorylated at Thr-174 can inhibit insulin secretion, BRSK2 is phosphorylated at Thr-260 can promote insulin secretion. It regulates reorganization of the actin cytoskeleton. It may play a role in the apoptotic response triggered by endoplasmatic reticulum (ER) stress.

Synonyms: C11orf7; FLJ41362; PEN11B; SAD1; STK29

Product images:



Host: Rabbit Target Name: BRSK2

Sample Type: HT1080 Whole Cell lysates

Antibody Dilution: 1.0ug/ml