

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA360210

# **RPLP0 Rabbit Polyclonal Antibody**

## **Product data:**

Product Type:	Primary Antibodies
Applications:	WB
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	The immunogen is a synthetic peptide directed towards the middle region of human RPLP0
Specificity:	Expected reactivity: Human
Formulation:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. Note that this product is shipped as lyophilized powder to China customers.
Concentration:	lot specific
Purification:	Affinity purified
Conjugation:	Unconjugated
Storage:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	34 kDa
Gene Name:	ribosomal protein lateral stalk subunit P0
Database Link:	<u>NP 000993.1</u>
	<u>Entrez Gene 6175 Human</u> <u>P05388</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### **GRIGENE** RPLP0 Rabbit Polyclonal Antibody – TA360210


Background:Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and<br/>a large 60S subunit. Together these subunits are composed of 4 RNA species and<br/>approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is<br/>a component of the 60S subunit. The protein, which is the functional equivalent of the E. coli<br/>L10 ribosomal protein, belongs to the L10P family of ribosomal proteins. It is a neutral<br/>phosphoprotein with a C-terminal end that is nearly identical to the C-terminal ends of the<br/>acidic ribosomal phosphoproteins P1 and P2. The P0 protein can interact with P1 and P2 to<br/>form a pentameric complex consisting of P1 and P2 dimers, and a P0 monomer. The protein<br/>is located in the cytoplasm. Transcript variants derived from alternative splicing exist; they<br/>encode the same protein. As is typical for genes encoding ribosomal proteins, there are<br/>multiple processed pseudogenes of this gene dispersed through the genome.

Synonyms: L10E; MGC88175; MGC111226; P0; PRLP0; RPP0

Ribosome

Protein Pathways:

### **Product images:**



Host: Rabbit Target Name: RPLP0 Sample Tissue: Human HCT116 Whole Cell Antibody Dilution: 1.0ug/ml

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US