

Product datasheet for TA361160

SKP2 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Reactivity: Human Host: Rabbit

Clonality: Polyclonal

Immunogen: The immunogen is a synthetic peptide directed towards the middle region of human SKP2

Expected reactivity: Human Specificity:

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

Concentration: lot specific

Purification: Affinity purified Conjugation: Unconjugated

For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small Storage:

aliquots to prevent freeze-thaw cycles.

Shelf life: one year from despatch. Stability:

Predicted Protein Size: 46 kDa

Gene Name: S-phase kinase-associated protein 2, E3 ubiquitin protein ligase

Database Link: NP 001230049.1

Entrez Gene 6502 Human

Q13309



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:

This gene encodes a member of the F-box protein family which is characterized by an approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbls class; in addition to an F-box, this protein contains 10 tandem leucine-rich repeats. This protein is an essential element of the cyclin A-CDK2 S-phase kinase. It specifically recognizes phosphorylated cyclin-dependent kinase inhibitor 1B (CDKN1B, also referred to as p27 or KIP1) predominantly in S phase and interacts with S-phase kinase-associated protein 1 (SKP1 or p19). In addition, this gene is established as a protooncogene causally involved in the pathogenesis of lymphomas. Alternative splicing of this gene generates three transcript variants encoding different isoforms.

Synonyms: FBL1; FBXL1; FLB1; MGC1366; p45skp2

Protein Families: Druggable Genome

Protein Pathways: Acute myeloid leukemia, Apoptosis, Cell cycle, Oocyte meiosis, p53 signaling pathway,

Pathways in cancer, Progesterone-mediated oocyte maturation, Small cell lung cancer,

Ubiquitin mediated proteolysis

Product images:

