

## **Product datasheet for TA361176**

# OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### **HLADQA1 (HLA-DQA1) Rabbit Polyclonal Antibody**

#### **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Reactivity: Human Host: Rabbit

**Clonality:** Polyclonal

Immunogen: The immunogen is a synthetic peptide directed towards the middle region of human HLA-

DQA1

**Specificity: Expected reactivity**: Human

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

**Concentration:** lot specific

Purification: Affinity purified
Conjugation: Unconjugated

**Storage:** For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small

aliquots to prevent freeze-thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 27 kDa

**Gene Name:** major histocompatibility complex, class II, DQ alpha 1

Database Link: NP 002113.2

Entrez Gene 3117 Human

P01909





Background:

HLA-DQA1 belongs to the HLA class II alpha chain paralogues. The class II molecule is a heterodimer consisting of an alpha (DQA) and a beta chain (DQB), both anchored in the membrane. It plays a central role in the immune system by presenting peptides derived from extracellular proteins. Class II molecules are expressed in antigen presenting cells. The alpha chain is approximately 33-35 kDa. It is encoded by 5 exons; exon 1 encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, and exon 4 encodes the transmembrane domain and the cytoplasmic tail. Within the DQ molecule both the alpha chain and the beta chain contain the polymorphisms specifying the peptide binding specificities, resulting in up to four different molecules. Typing for these polymorphisms is routinely done for bone marrow transplantation

Synonyms: CD; CELIAC1; DQ-A1; GSE; HLA-DQA

**Protein Families:** Transmembrane

**Protein Pathways:** Allograft rejection, Antigen processing and presentation, Asthma, Autoimmune thyroid

disease, Cell adhesion molecules (CAMs), Graft-versus-host disease, Systemic lupus

erythematosus, Type I diabetes mellitus, Viral myocarditis

#### **Product images:**

