

### **Product datasheet for TA361266**

#### 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

## **CCN4 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WB

Reactivity: Human Rabbit

Clonality: Polyclonal

**Immunogen:** The immunogen is a synthetic peptide directed towards the middle region of human WISP1

**Specificity: Expected reactivity**: Human

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

**Concentration:** lot specific

Purification: Affinity purified
Conjugation: Unconjugated

Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small

aliquots to prevent freeze-thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 40 kDa

**Gene Name:** WNT1 inducible signaling pathway protein 1

Database Link: NP 001191798.1

Entrez Gene 8840 Human

<u>095388</u>




#### Background:

This gene encodes a member of the WNT1 inducible signaling pathway (WISP) protein subfamily, which belongs to the connective tissue growth factor (CTGF) family. WNT1 is a member of a family of cysteine-rich, glycosylated signaling proteins that mediate diverse developmental processes. The CTGF family members are characterized by four conserved cysteine-rich domains: insulin-like growth factor-binding domain, von Willebrand factor type C module, thrombospondin domain and C-terminal cystine knot-like domain. This gene may be downstream in the WNT1 signaling pathway that is relevant to malignant transformation. It is expressed at a high level in fibroblast cells, and overexpressed in colon tumors. The encoded protein binds to decorin and biglycan, two members of a family of small leucine-rich proteoglycans present in the extracellular matrix of connective tissue, and possibly prevents the inhibitory activity of decorin and biglycan in tumor cell proliferation. It also attenuates p53-mediated apoptosis in response to DNA damage through activation of the Akt kinase. It is 83% identical to the mouse protein at the amino acid level. Multiple alternatively spliced transcript variants have been identified.

Synonyms: CCN4; WISP-1; WISP1c; WISP1tc

# **Product images:**



Host: Rabbit Target Name: WISP1

Sample Tissue: Human MCF7 Whole Cell lysates

Antibody Dilution: 1ug/ml