

## Product datasheet for **TA361423**

### SUCLA2 Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	WB
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	The immunogen is a synthetic peptide directed towards the middle terminal region of human SUCLA2
Specificity:	<b>Expected reactivity:</b> Human
Formulation:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. <i>Note that this product is shipped as lyophilized powder to China customers.</i>
Concentration:	lot specific
Purification:	Affinity purified
Conjugation:	Unconjugated
Storage:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	48 kDa
Gene Name:	succinate-CoA ligase ADP-forming beta subunit
Database Link:	<a href="#">NP_003841.1</a> <a href="#">Entrez Gene 8803 Human</a> <a href="#">Q9P2R7-2</a>



[View online »](#)

**Background:**


Succinyl-CoA synthetase (SCS) is a mitochondrial matrix enzyme that acts as a heterodimer, being composed of an invariant alpha subunit and a substrate-specific beta subunit. The protein encoded by this gene is an ATP-specific SCS beta subunit that dimerizes with the SCS alpha subunit to form SCS-A, an essential component of the tricarboxylic acid cycle. SCS-A hydrolyzes ATP to convert succinate to succinyl-CoA. Defects in this gene are a cause of myopathic mitochondrial DNA depletion syndrome. A pseudogene of this gene has been found on chromosome 6.

**Synonyms:**

A-BETA; SCS-betaA

**Protein Pathways:**

Citrate cycle (TCA cycle), Metabolic pathways, Propanoate metabolism

**Product images:**

Host: Rabbit  
Target Name: SUCLA2  
Sample Type: Testicular Tumor Lysate  
Antibody Dilution: 1.0µg/ml

Host: Rabbit  
Target Name: SUCLA2  
Sample Tissue: Human Testicular Tumor lysates  
Antibody Dilution: 1ug/ml