

## Product datasheet for **TA361897**

### Thymine DNA glycosylase (TDG) Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	WB
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	The immunogen is a synthetic peptide directed towards the middle terminal region of human TDG
Specificity:	<b>Expected reactivity:</b> Human
Formulation:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. <i>Note that this product is shipped as lyophilized powder to China customers.</i>
Concentration:	lot specific
Purification:	Affinity purified
Conjugation:	Unconjugated
Storage:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	45 kDa
Gene Name:	thymine DNA glycosylase
Database Link:	<a href="#">NP_003202.3</a> <a href="#">Entrez Gene 6996 Human Q13569</a>



[View online »](#)

**Background:**

The protein encoded by this gene belongs to the TDG/mug DNA glycosylase family. Thymine-DNA glycosylase (TDG) removes thymine moieties from G/T mismatches by hydrolyzing the carbon-nitrogen bond between the sugar-phosphate backbone of DNA and the mispaired thymine. With lower activity, this enzyme also removes thymine from C/T and T/T mispairings. TDG can also remove uracil and 5-bromouracil from mispairings with guanine. This enzyme plays a central role in cellular defense against genetic mutation caused by the spontaneous deamination of 5-methylcytosine and cytosine. This gene may have a pseudogene in the p arm of chromosome 12.

**Synonyms:**


E130317C12Rik; hTDG; JZA-3; Jza1; OTTMUSP00000028912; OTTMUSP00000028913

**Protein Families:**

Druggable Genome

**Protein Pathways:**

Base excision repair

**Product images:**

Host: Rabbit  
Target Name: TDG  
Sample Type: HCT116 Cell Lysate  
Antibody Dilution: 1.0µg/ml

Host: Rabbit  
Target Name: TDG  
Sample Tissue: Human HCT116 Whole Cell lysates  
Antibody Dilution: 1 ug/ml