

Product datasheet for TA362377

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Proteasome Activator Subunit 4 (PSME4) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Reactivity: Human Host: Rabbit

Clonality: Polyclonal

Immunogen: The immunogen is a synthetic peptide directed towards the C terminal region of human

PSME4

Specificity: Expected reactivity: Human

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

Concentration: lot specific

Purification: Affinity purified
Conjugation: Unconjugated

Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small

aliquots to prevent freeze-thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 133 kDa

Gene Name: proteasome activator subunit 4

Database Link: NP 055429.2

Entrez Gene 23198 Human

Q14997-2






Background:

Associated component of the proteasome that specifically recognizes acetylated histones and promotes ATP- and ubiquitin-independent degradation of core histones during spermatogenesis and DNA damage response. Recognizes and binds acetylated histones via its bromodomain-like (BRDL) region and activates the proteasome by opening the gated channel for substrate entry. Binds to the core proteasome via its C-terminus, which occupies the same binding sites as the proteasomal ATPases, opening the closed structure of the proteasome via an active gating mechanism. Component of the spermatoproteasome, a form of the proteasome specifically found in testis: binds to acetylated histones and promotes degradation of histones, thereby participating actively to the exchange of histones during spermatogenesis. Also involved in DNA damage response in somatic cells, by promoting degradation of histones following DNA double-strand breaks.

Synonyms: FLJ21864; KIAA0077; MGC138374; MGC142228; PA200

Protein Pathways: Proteasome

Product images:



Host: Rabbit Target Name: PSME4

Sample Tissue: Human HCT116 Whole Cell

lysates

Antibody Dilution: 1ug/ml