

Product datasheet for TA362972

Rbpj Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Reactivity: Mouse Host: Rabbit

Clonality: Polyclonal

Immunogen: The immunogen is a synthetic peptide directed towards the middle region of mouse RBPI

Expected reactivity: Mouse Specificity:

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

Concentration: lot specific

Purification: Affinity purified Conjugation: Unconjugated

For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small Storage:

aliquots to prevent freeze-thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 53 kDa

Gene Name: recombination signal binding protein for immunoglobulin kappa J region

Database Link: NP 001074396.1

Entrez Gene 19664 Mouse

P31266-2



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Transcriptional regulator that plays a central role in Notch signaling, a signaling pathway involved in cell-cell communication that regulates a broad spectrum of cell-fate determinations. Acts as a transcriptional repressor when it is not associated with Notch proteins. When associated with some NICD product of Notch proteins (Notch intracellular domain), it acts as a transcriptional activator that activates transcription of Notch target genes. Probably represses or activates transcription via the recruitment of chromatin remodeling complexes containing histone deacetylase or histone acetylase proteins, respectively. Specifically binds to the immunoglobulin kappa-type J segment recombination signal sequence. Binds specifically to methylated DNA. Binds to the oxygen responsive element of COX4I2 and activates its transcription under hypoxia conditions (4% oxygen) (By similarity). Negatively regulates the phagocyte oxidative burst in response to bacterial infection by repressing transcription of NADPH oxidase subunits.

Synonyms:

CBF-1; CBF1; csl; IGKJRB; IGKJRB1; KBF2; MGC61669; OTTHUMP00000123441; RBP-J; RBP-JK; RBPJK; RBPSUH; SUH

Product images:



Host: Rabbit Target Name: RBPJ

Sample Tissue: Mouse Heart lysates Antibody Dilution: 1ug/ml