

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA363032

Fzd4 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	The immunogen is a synthetic peptide directed towards the C terminal region of mouse FZD4
Specificity:	Expected reactivity: Mouse
Formulation:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. Note that this product is shipped as lyophilized powder to China customers.
Concentration:	lot specific
Purification:	Affinity purified
Conjugation:	Unconjugated
Storage:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	60 kDa
Gene Name:	frizzled class receptor 4
Database Link:	<u>NP_032081.3</u> <u>Entrez Gene 14366 Mouse</u> <u>Q61088</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE Fzd4 Rabbit Polyclonal Antibody – TA363032

Background:Receptor for Wnt proteins. Most of frizzled receptors are coupled to the beta-catenin
(CTNNB1) canonical signaling pathway, which leads to the activation of disheveled proteins,
inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin (CTNNB1) and activation of
Wnt target genes. Plays a critical role in retinal vascularization by acting as a receptor for Wnt
proteins and norrin (NDP). In retina, it can be both activated by Wnt protein-binding, but also
by a Wnt-independent signaling via binding of norrin (NDP), promoting in both cases beta-
catenin (CTNNB1) accumulation and stimulation of LEF/TCF-mediated transcriptional
programs. A second signaling pathway involving PKC and calcium fluxes has been seen for
some family members, but it is not yet clear if it represents a distinct pathway or if it can be
integrated in the canonical pathway, as PKC seems to be required for Wnt-mediated
inactivation of GSK-3 kinase. Both pathways seem to involve interactions with G-proteins. May
be involved in transduction and intercellular transmission of polarity information during
tissue morphogenesis and/or in differentiated tissues. Activation by Wnt5A stimulates PKC
activity via a G-protein-dependent mechanism.

Synonyms: CD344; EVR1; FEVR; Fz-4; FZD4S; FzE4; GPCR; hFz4; MGC34390

Product images:



Host: Rabbit Target Name: FZD4 Sample Tissue: Mouse Testis lysates Antibody Dilution: 1ug/ml

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US