

## **Product datasheet for TA363706**

# Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### **CCN4 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WB

Reactivity: Human Rabbit

Clonality: Polyclonal

**Immunogen:** The immunogen is a synthetic peptide directed towards the C terminal region of human

WISP1

**Specificity: Expected reactivity**: Human

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

**Concentration:** lot specific

Purification: Affinity purified
Conjugation: Unconjugated

**Storage:** For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small

aliquots to prevent freeze-thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 21 kDa

**Gene Name:** WNT1 inducible signaling pathway protein 1

**Database Link:** <u>NP 001191798.1</u>

Entrez Gene 8840 Human

O95388-5






#### Background:

This gene encodes a member of the WNT1 inducible signaling pathway (WISP) protein subfamily, which belongs to the connective tissue growth factor (CTGF) family. WNT1 is a member of a family of cysteine-rich, glycosylated signaling proteins that mediate diverse developmental processes. The CTGF family members are characterized by four conserved cysteine-rich domains: insulin-like growth factor-binding domain, von Willebrand factor type C module, thrombospondin domain and C-terminal cystine knot-like domain. This gene may be downstream in the WNT1 signaling pathway that is relevant to malignant transformation. It is expressed at a high level in fibroblast cells, and overexpressed in colon tumors. The encoded protein binds to decorin and biglycan, two members of a family of small leucine-rich proteoglycans present in the extracellular matrix of connective tissue, and possibly prevents the inhibitory activity of decorin and biglycan in tumor cell proliferation. It also attenuates p53-mediated apoptosis in response to DNA damage through activation of the Akt kinase. It is 83% identical to the mouse protein at the amino acid level. Multiple alternatively spliced transcript variants have been identified.

Synonyms: CCN4; WISP-1; WISP1c; WISP1tc

#### **Product images:**



Host: Rabbit Target Name: WISP1

Sample Tissue: Human Jurkat Whole Cell lysates

Antibody Dilution: 1ug/ml