

Product datasheet for TA363856

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD11b/c Mouse Monoclonal Antibody [Clone ID: OX-42]

Product data:

Product Type: Primary Antibodies

Clone Name: OX-42
Applications: IHC

Recommended Dilution: Tested for immunohistochemistry (IHC); has been described to work in FACS. Approximate

working dilution for IHC: Frozen sections: 5 -10µg/ml (1:100 - 1:200) Paraffin sections: does not react on routinely processed paraffin sections. Optimal dilutions should be determined

by the end user. Suggested positive control: Rat spleen.

Reactivity: Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Peritoneal macrophages.

Specificity: Rat: granulocytes, macrophages, microglial cells. Other species: not tested

Epitope: CD11b/c. OX-42 recognises three major polypeptides of 160kDa, 103kDa and 95kDa and a minor 133kDa protein under non-reducing conditions. Under reducing conditions two

major peptides of 163kDa and 100kDa and a minor 135kDa peptide are detected.

Distribution: Isolated cells: OX-42 shows a similar reaction pattern to OX- 41 on various isolated cells but stains bronchial lavage cells to a lesser degree. OX-42 inhibits complement mediated rosettes. Tissue sections: OX-42 is a most suitable marker for the identification of microglial cells and Kupffer cells. It also stains dendritic cells strongly. Only a subpopulation

of alveolar macrophages is recognised.

Formulation: Purified, liquid Supplied as 1ml solution. This stock solution contains 1mg/ml IgG, phosphate

buffered saline pH 7.2 (PBS) and 0.09% sodium azide as a preservative.

Concentration: lot specific

Conjugation: Unconjugated


Storage: Stock solution or aliquots thereof: 1 year at -20°C. Avoid repeated thawing and freezing.



Background:

Monoclonal antibody OX-42 recognises most macrophages (including resident peritoneal and activated macrophages), Kupffer cells, but only about 35% of alveolar macrophages. It also labels dendritic cells extensively, and granulocytes and cells with the morphology of microglia in the brain. OX-42 is probably the rat equivalent of the human CR3 complement receptor for iC3b. In combination with OX-41 it facilitates a more distinct phenotyping of resident macrophages.

Product images:



TA363856, Clone OX-42, rat spleen, frozen section