

Product datasheet for TA364013

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Bombesin Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA

Recommended Dilution: This antibody has been tested and validated in ELISA against Bombesin. Other applications

like immunohistochemistry (IHC), FACS or Western Blot may work as well. Optimal dilutions

should be determined by the end user.

Reactivity: Human, Mammalian

Host: Rabbit

Clonality: Polyclonal

Immunogen: Synthetic peptide Pyr-Gln-Arg-Leu-Gly-Asn-Gln-Trp-Ala-Val-Gly-His- Leu-Met-NH2 coupled to

carrier protein.

Formulation: Each vial contains enough antiserum for 500 RIA tubes. The powder should be rehydrated

with 50ml RIA buffer. Upon reconstitution to 50ml total volume, the solution contains 0.1M sodium phosphate buffer (pH 7.4), 0.05M NaCl, 0.1% BSA, 0.01% NaN3, and 0.1% Triton X-100. Store at 4°C. This should ensure antibody stability for approximately one month.

Concentration: N/A

Conjugation: Unconjugated

Storage: Original vial: at least one year at 4° - 8°C from date of delivery. Minimize repeated thawing

and freezing of the antiserum by freezing aliquots at -20°C or below.

Background: Bombesin is a 14-amino acid peptide originally isolated from the skin of the European fire-

bellied toad (Bombina bombina). It has two known homologs in mammals called neuromedin

B and gastrin-releasing peptide. Bombesin stimulates gastrin release from G cells and activates three different G-protein-coupled receptors known as BBR1, -2, and -3. It also

activates these receptors in the brain. Together with cholecystokinin, Bombesin is the second major source of negative feedback signals that stop eating behaviour. Also, it is known to be a tumor marker for small cell carcinoma of lung, gastric cancer, pancreatic cancer, and neuroblastoma. This antibody was generated by immunization of rabbits with Bombesin

coupled to a carrier protein.

