

Product datasheet for TA364323

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Vasopressin (AVP) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA

Recommended Dilution: This antibody has been tested and validated in ELISA against Desmopressin. Other

applications like immunohistochemistry (IHC), FACS or Western Blot may work as well.

Optimal dilutions should be determined by the end user.

Reactivity: Human, Mammalian

Host: Rabbit

Clonality: Polyclonal

Immunogen: Synthetic peptide 3-Mercaptopropionyl-Tyr-Phe-Gln-Asn-Cys-Pro-D-Arg- Gly-NH2, (Disulfide

bond) coupled to carrier protein.

Formulation: Each vial contains enough antiserum for 500 RIA tubes. The powder should be rehydrated

with 50ml RIA buffer. Upon reconstitution to 50ml total volume, the solution contains 0.1M sodium phosphate buffer (pH 7.4), 0.05M NaCl, 0.1% BSA, 0.01% NaN3, and 0.1% Triton X-100. Store at 4°C. This should ensure antibody stability for approximately one month.

Concentration: N/A

Conjugation: Unconjugated

Storage: Original vial: at least one year at 4° - 8°C from date of delivery. Minimize repeated thawing

and freezing of the antiserum by freezing aliquots at -20°C or below.

Gene Name: arginine vasopressin

Database Link: Entrez Gene 551 Human

P01185



Vasopressin (AVP) Rabbit Polyclonal Antibody - TA364323

Background: Desmopressin is a synthetic octapeptide, analogue of human hormone arginine vasopressin,

with antidiuretic and coagulant activities. Desmopressin binds to V2 receptors in renal collecting ducts which leads to exocytosis of von Willebrand factor (VWF) and tissue plasminogen activator (t-PA) from Weibel-Palade bodies, thereby increasing water resorption.

This agent also increases nitric oxide (NO) production via activation of endothelial NO synthase, thereby induces afferent arteriolar vasodilation. Furthermore, desmopressin stimulates the release of factor VIII from endothelial cells mediated through V1a receptor,

thereby promotes blood coagulation. This antibody was generated by immunization of

rabbits with Desmopressin coupled to a carrier protein.

Synonyms: ADH; ARVP; AVP-NPII; AVRP; neurohypophyseal; VP