

Product datasheet for TA364442

COX4NB (EMC8) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: A549, Hela and HT-29 cells

IHC: 50-200

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human EMC8

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 24 kDa

Gene Name: ER membrane protein complex subunit 8

Database Link: Entrez Gene 10328 Human

043402

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn






Background:


COX4NB (Neighbor of COX4) is a 210 amino acid protein encoded by the human gene COX4NB. COX4NB belongs to the UPF0172 (NOC4) family and is found on chromosome 16, adjacent to the gene that encodes COX4. The GAN gene is located on chromosome 16 and, with mutation, may lead to giant axonal neuropathy, a nervous system disorder characterized by increasing malfunction with growth. The rare disorder Rubinstein-Taybi syndrome is also associated with chromosome 16 through the CREBBP gene which encodes a critical CREB binding protein. Crohn's disease is a gastrointestinal inflammatory condition associated with chromosome 16 through the NOD2 gene. An association with systemic lupus erythematosis and a number of other auto-immune disorders with the pericentromeric region of chromosome 16 has led to the identification of SLC5A11 as a potential autoimmune modifier.

Synonyms: Cox4nb; Fam158b; Noc4

Product images:




Gel: 10%SDS-PAGE
Lysate: 40 µg
Lane 1-3: A549 cells
Hela cells
HT29 cells
Primary antibody: TA364442 (EMC8 Antibody) at dilution 1/600
Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution
Exposure time: 1 minute




Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA364442 (EMC8 Antibody) at dilution 1/30 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA364442 (EMC8 Antibody) at dilution 1/30, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA364442 (EMC8 Antibody) at dilution 1/30 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA364442 (EMC8 Antibody) at dilution 1/30, treated with fusion protein. (Original magnification: ×200)