

Product datasheet for TA364622S

IDH3G Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB: 200-1000 WB positive control: A431 cells and mouse brain tissue IHC: 25-100 Positive control: Human thyroid cancer Predicted cell location: Cytoplasm
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Immunogen:	Fusion protein of human IDH3G
Formulation:	pH7.4 PBS, 0.05% NaN3, 40% Glycerol
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C.
Stability:	1 year
Predicted Protein Size:	43 kDa
Gene Name:	isocitrate dehydrogenase 3 (NAD(+)) gamma
Database Link:	<u>Entrez Gene 3421 Human</u> <u>P51553</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

DRIGENE IDH3G Rabbit Polyclonal Antibody – TA364622S


Background: Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. NAD(+)-dependent isocitrate dehydrogenases catalyze the allosterically regulated rate-limiting step of the tricarboxylic acid cycle. Each isozyme is a heterotetramer that is composed of two alpha subunits, one beta subunit, and one gamma subunit.

Synonyms: H-IDHG; IDH-gamma

Product images:




Gel: 10%SDS-PAGE Lysate: 40 µg Lane 1-2: A431 cells mouse brain tissue Primary antibody: [TA364622] (IDH3G Antibody) at dilution 1/300 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 15 seconds



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA364622] (IDH3G Antibody) at dilution 1/30 (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA364622] (IDH3G Antibody) at dilution 1/30, treated with fusion protein. (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA364622] (IDH3G Antibody) at dilution 1/30 (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA364622] (IDH3G Antibody) at dilution 1/30, treated with fusion protein. (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US