

## **Product datasheet for TA364635**

# **PITX2 Rabbit Polyclonal Antibody**

### **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB: 500-2000

WB positive control: HepG2 cells

Reactivity: Human, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Fusion protein of human PITX2

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 35 kDa

Gene Name: paired like homeodomain 2

Database Link: Entrez Gene 5308 Human

Q99697

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn






Background:

This gene encodes a member of the RIEG/PITX homeobox family, which is in the bicoid class of homeodomain proteins. The encoded protein acts as a transcription factor and regulates procollagen lysyl hydroxylase gene expression. This protein plays a role in the terminal differentiation of somatotroph and lactotroph cell phenotypes, is involved in the development of the eye, tooth and abdominal organs, and acts as a transcriptional regulator involved in basal and hormone-regulated activity of prolactin. Mutations in this gene are associated with Axenfeld-Rieger syndrome, iridogoniodysgenesis syndrome, and sporadic cases of Peters anomaly. A similar protein in other vertebrates is involved in the determination of left-right asymmetry during development. Alternatively spliced transcript variants encoding distinct isoforms have been described.

Synonyms:

ARP1; Brx1; IDG2; IGDS; IGDS2; IHG2; IRID2; MGC20144; MGC111022; Otlx2; OTTHUMP00000163736; PTX2; RGS; RIEG; RIEG1; RS; Solurshin

## **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 µg Lane: HepG2 cells Primary antibody: TA364635 (PITX2 Antibody) at dilution 1/200 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution