

# Product datasheet for TA364707

# IL2 Receptor beta (IL2RB) Rabbit Polyclonal Antibody

# **Product data:**

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB: 500-2000 WB positive control: 231, 293T, Raji and hela cells IHC: 25-100 Positive control: Human thyroid cancer Predicted cell location: Cytoplasm
Reactivity:	Human
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Immunogen:	Fusion protein of human IL2RB
Formulation:	pH7.4 PBS, 0.05% NaN3, 40% Glycerol
Concentration:	lot specific
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C.
Stability:	1 year
Predicted Protein Size:	61 kDa
Gene Name:	interleukin 2 receptor subunit beta
Database Link:	<u>Entrez Gene 3560 Human</u> <u>P14784</u>

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

# Second Se


## Background: The interleukin 2 receptor, which is involved in T cell-mediated immune responses, is present in 3 forms with respect to ability to bind interleukin 2. The low affinity form is a monomer of the alpha subunit and is not involved in signal transduction. The intermediate affinity form consists of an alpha/beta subunit heterodimer, while the high affinity form consists of an alpha/beta/gamma subunit heterotrimer. Both the intermediate and high affinity forms of the receptor are involved in receptor-mediated endocytosis and transduction of mitogenic signals from interleukin 2. The protein encoded by this gene represents the beta subunit and is a type I membrane protein.

**Synonyms:** CD122; P70-75; p75

## **Product images:**




Gel: 10%SDS-PAGE Lysate: 40 µg Lane 1-4: 231 cells 293T cells Raji cells hela cells Primary antibody: TA364707 (IL2RB Antibody) at dilution 1/325 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 20 seconds



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA364707 (IL2RB Antibody) at dilution 1/30 (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA364707 (IL2RB Antibody) at dilution 1/30, treated with fusion protein. (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA364707 (IL2RB Antibody) at dilution 1/30 (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA364707 (IL2RB Antibody) at dilution 1/30, treated with fusion protein. (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US