

Product datasheet for TA364930S

PPM1A Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 1000-5000

WB positive control: Hela cell lysate

IHC: 150-300

Positive control: Human cervical cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human PPM1A

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 42 kDa

Gene Name: protein phosphatase, Mg2+/Mn2+ dependent 1A

Database Link: Entrez Gene 5494 Human

P35813

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn






Background:

The protein encoded by this gene is a member of the PP2C family of Ser/Thr protein phosphatases. PP2C family members are known to be negative regulators of cell stress response pathways. This phosphatase dephosphorylates, and negatively regulates the activities of, MAP kinases and MAP kinase kinases. It has been shown to inhibit the activation of p38 and JNK kinase cascades induced by environmental stresses. This phosphatase can also dephosphorylate cyclin-dependent kinases, and thus may be involved in cell cycle control. Overexpression of this phosphatase is reported to activate the expression of the tumor suppressor gene TP53/p53, which leads to G2/M cell cycle arrest and apoptosis. Three alternatively spliced transcript variants encoding distinct isoforms have been described. [provided by RefSeq, Jul 2008]

Synonyms:

FLJ42306; IA; MGC9201; PP2C-ALPHA; PP2CA; PP2Calpha; PPPM1A

Product images:



Gel: 8%SDS-PAGE Lysate: 40 µg Lane: Hela cell lysate


Primary antibody: [TA364930] (PPM1A Antibody)

at dilution 1/1000

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 10 seconds




Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA364930] (PPM1A Antibody) at dilution 1/200 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA364930] (PPM1A Antibody) at dilution 1/200, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA364930] (PPM1A Antibody) at dilution 1/200 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA364930] (PPM1A Antibody) at dilution 1/200, treated with fusion protein. (Original magnification: ×200)