

## **Product datasheet for TA364995S**

## **SAA4 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Human normal stomach tissue lysate

IHC: 25-100

Positive control: Human cervical cancer

Predicted cell location: Secreted

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Full length fusion protein

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 15 kDa

**Gene Name:** serum amyloid A4, constitutive

Database Link: Entrez Gene 6291 Human

P35542



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com






Background:

SAA4 is a member of the SAA family.SAA proteins are family of apolipoproteins of high density lipoprotein (HDL). They can be separated into two distinct groups. First group (SAA1, SAA2, and SAA3) consists of acute phase reactant whose expression level increase in the blood in a response to trauma, infection, inflammation, and neoplasia. These acute phase SAAs associates with HDL during inflammation and remodel the HDL particle by displacing Apo-A1. The second distinct group consists of SAA4 and SAA5 which exist as the minor apolipoproteins on HDL, but this group of SAA constitutes more than 90% of all the SAA during homeostasis, and it is thought to play a role in the normal functioning of the HDL particle. SAA4 is a constitutively expressed protein which expressed only in humans and mice.It is connected almost completely with lipoproteins of the high density range. The physiological function of SAA4 is unknown, and its serum concentration has no association with those of other major apolipoproteins.

Synonyms: C-SAA; CSAA

## **Product images:**



Gel: 12%SDS-PAGE Lysate: 40 μg

Lane: Human normal stomach tissue lysate Primary antibody: [TA364995] (SAA4 Antibody) at


dilution 1/350

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 5 minutes





Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA364995] (SAA4 Antibody) at dilution 1/35 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA364995] (SAA4 Antibody) at dilution 1/35, treated with fusion protein. (Original magnification: ×200)