

# Product datasheet for TA365373S

**Primary Antibodies** 

WB positive control: A549 cell

Fusion protein of human ZNF281

Antigen affinity purification

zinc finger protein 281

Entrez Gene 23528 Human

pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Positive control: Human thyroid cancer

Predicted cell location: Cytoplasm and Nucleus

IHC, WB

IHC: 60-450

Human, Mouse

Rabbit

Polyclonal

Unconjugated Store at -20°C.

1 year

97 kDa

Q9Y2X9

lgG

# **ZNF281 Rabbit Polyclonal Antibody**

Recommended Dilution: WB: 500-2000

# **Product data:**

**Product Type:** 

**Applications:** 

**Reactivity:** 

Host:

Isotype:

**Clonality:** 

Immunogen:

Formulation:

**Purification:** 

**Conjugation:** 

Gene Name:

Database Link:

**Predicted Protein Size:** 

Storage: Stability:

#### **OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

|--|

View online »


This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

## **GRIGENE** ZNF281 Rabbit Polyclonal Antibody – TA365373S


**Background:** Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. ZNF281, also known as GC-box-binding zinc finger protein 1, ZBP-99 or ZNP-99 (zinc finger DNA-binding protein 99), is a zinc finger protein that belongs to the Krüppel C2H2-type zinc finger protein family. It is expressed ubiquitously at low levels with predominant expression in kidney, liver, lymphocytes and placenta. ZNF281 localizes to the nucleus and contains four C2H2-type zinc fingers. ZNF281 plays a role in repressing the transcription of a variety of genes including Gastrin and ODC (ornithine decarboxylase). In particular, ZNF281 functions by binding to the G-rich box in the enhancer region of the gene. Upon DNA damage, ZNF281 may become phosphorylated by Atm or ATR.

Synonyms: FLJ12859; FLJ14378; GZP1; OTTHUMP00000033838; ZBP-99; ZBP99; ZNP-99

### **Product images:**




Gel: 6%SDS-PAGE Lysate: 40 µg Lane: A549 cell Primary antibody: [TA365373] (ZNF281 Antibody) at dilution 1/600 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 5 seconds



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA365373] (ZNF281 Antibody) at dilution 1/70 (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA365373] (ZNF281 Antibody) at dilution 1/70, treated with fusion protein. (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human brain tissue using [TA365373] (ZNF281 Antibody) at dilution 1/70 (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human brain tissue using [TA365373] (ZNF281 Antibody) at dilution 1/70, treated with fusion protein. (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US