

Product datasheet for TA365439S

VDAC1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 1000-5000

WB positive control: Jurkat cell, 293T cell, Mouse kidney tissue lysates

IHC: 150-300

Positive control: Human prostate cancer

Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human VDAC1

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 31 kDa

Gene Name: voltage dependent anion channel 1

Database Link: Entrez Gene 7416 Human

P21796



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Voltage-dependent anion channel (VDAC), ubiquitously expressed and located in the outer mitochondrial membrane, is generally thought to be the primary means by which metabolites diffuse in and out of the mitochondria. In addition, this channel plays a role in apoptotic signaling. The change in mitochondrial permeability characteristic of apoptosis is mediated by Bcl-2 family proteins, which bind to VDAC, altering the channel kinetics. Homodimerization of VDAC may be a mechanism for changing mitochondrial permeability and supporting release of cytochrome c. In mammalian cells, there are three VDAC isoforms, VDAC1, which is the most widely expressed isoform, as well as VDAC2 and VDAC3.

Synonyms:


hVDAC1; MGC111064; OTTHUMP00000159381; OTTHUMP00000165946; PORIN; PORIN-31-HL; VDAC; VDAC-1

Product images:




Gel: 12%SDS-PAGE Lysate: 40 µg Lane 1-3: Jurkat cell 293T cell Mouse kidney tissue lysates Primary antibody: [TA365439] (VDAC1 Antibody) at dilution 1/1150 Secondary antibody: Goat anti rabbit IgG at 1/5000 dilution

Exposure time: 30 seconds




Immunohistochemistry of paraffin-embedded Human prostate cancer tissue using [TA365439] (VDAC1 Antibody) at dilution 1/150 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human prostate cancer tissue using [TA365439] (VDAC1 Antibody) at dilution 1/150, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA365439] (VDAC1 Antibody) at dilution 1/150 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA365439] (VDAC1 Antibody) at dilution 1/150, treated with fusion protein. (Original magnification: ×200)