

Product datasheet for TA365517S

BDH2 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 500-2000

WB positive control: 293T and A172 cell

Mouse kidney tissue lysates

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Full length fusion protein

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 27 kDa

Gene Name: 3-hydroxybutyrate dehydrogenase, type 2

Database Link: Entrez Gene 56898 Human

Q9BUT1



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

DHRS6 (dehydrogenase/reductase SDR family member 6), also known as EFA6R, SDR15C1, UCPA-OR,UNQ6308 or BDH2, is a 245 amino acid cytoplasmic protein belonging to the short-chain dehydrogenases/reductases (SDR) family, an evolutionarily conserved family of oxidoreductases found in all forms of life. DHRS6 is a novel, cytosolic type II R- β -hydroxybutyrate dehydrogenase that exists as two alternatively spliced isoforms and may have an essential role as a nutrient or building block in cellular survival. Human DHRS6 and its vertebrate orthologs show high levels of sequence identities to bacterial hydroxybutyrate dehydrogenases. DHRS6 may play an important role in the peripheral utilization of 3-hydroxybutyrate and its cytoplasmic localization with its high ratio of oxidized NAD+, the NAD+ dependence and the kinetic parameters of DHRS6 make it suitable to convert high levels of circulating 3-hydroxybutyrate into acetoacetate.

Synonyms:

DHRS6; EFA6R; FLJ13261; PRO20933; SDR15C1; UCPA-OR; UNQ6308

Product images:



Gel: 12%SDS-PAGE Lysate: 40 μg

Lane 1-3:293T and A172 cell Mouse kidney tissue lysates

Primary antibody: [TA365517] (BDH2 Antibody) at

dilution 1/300

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 15 seconds