

# **Product datasheet for TA365543S**

## C14orf130 (UBR7) Rabbit Polyclonal Antibody

#### **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: Hela cell lysate

IHC: 30-150

Positive control: Human colorectal cancer

Predicted cell location: Nucleus

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Fusion protein of human UBR7

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 48 kDa

**Gene Name:** ubiquitin protein ligase E3 component n-recognin 7 (putative)

Database Link: Entrez Gene 55148 Human

Q8N806



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




#### Background:

Ubiquitination is an important mechanism through which three classes of enzymes act in concert to target short-lived or abnormal proteins for destruction. The three classes of enzymes involved in ubiquitination are the ubiquitin-activating enzymes (E1s), the ubiquitin-conjugating enzymes (E2s) and the ubiquitin-protein ligases (E3s). Ubr7 (ubiquitin protein ligase E3 component n-recognin 7), also known as C14orf130 or N-recognin-7, is a 425 amino acid protein that contains one UBR-type zinc finger and one PHD zinc finger. Participating in protein modification events within the N-end rule pathway, Ubr7 functions as an E3 ubiquitin-protein ligase that recognizes and binds proteins that contain destabilizing N-terminal residues, thereby leading to their ubiquitination and subsequent degradation.

Synonyms:

C14orf130; FLJ10483; MGC9518; N-recognin-7

### **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 µg Lane: Hela cell lysate


Primary antibody: [TA365543] (UBR7 Antibody) at

dilution 1/300


Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution


Exposure time: 1 minute




Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using [TA365543] (UBR7 Antibody) at dilution 1/30 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using [TA365543] (UBR7 Antibody) at dilution 1/30, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA365543] (UBR7 Antibody) at dilution 1/30 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA365543] (UBR7 Antibody) at dilution 1/30, treated with fusion protein. (Original magnification: ×200)