

Product datasheet for TA365593S

CCDC112 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: 231 cell lysate

IHC: 25-100

Positive control: Human brain

Predicted cell location: Cytoplasm or Nucleus

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human CCDC112

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 54 kDa

Gene Name: coiled-coil domain containing 112

Database Link: Entrez Gene 153733 Human

Q8NEF3

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn






Background:

CCDC112 (coiled-coil domain containing 112), also known as MBC1 (mutated in bladder cancer 1), is a 446 amino acid protein. The gene encoding CCDC112 is located on chromosome 5. Due to alternative splicing events, CCDC112 exists as two isoforms. Chromosome 5 comprises about 6% of human genomic DNA and contains 181 million base pairs encoding around 1,000 genes. It is associated with Cockayne syndrome through the ERCC8 gene and familial adenomatous polyposis through the adenomatous polyposis coli (APC) tumor suppressor gene. Treacher Collins syndrome is also chromosome 5 associated and is caused by insertions or deletions within the TCOF1 gene. Deletion of the p arm of chromosome 5 leads to Cri du chat syndrome. Deletion of 5q or chromosome 5 altogether is common in therapy-related acute myelogenous leukemias and myelodysplastic syndrome.

Synonyms: MBC1; MGC39633

Product images:



Gel: 8%SDS-PAGE Lysate: 40 µg Lane: 231 cell lysate


Primary antibody: [TA365593] (CCDC112

Antibody) at dilution 1/500

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 15 seconds




Immunohistochemistry of paraffin-embedded Human brain tissue using [TA365593] (CCDC112 Antibody) at dilution 1/30 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human brain tissue using [TA365593] (CCDC112 Antibody) at dilution 1/30, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human prostate cancer tissue using [TA365593] (CCDC112 Antibody) at dilution 1/30 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human prostate cancer tissue using [TA365593] (CCDC112 Antibody) at dilution 1/30, treated with fusion protein. (Original magnification: ×200)