

Product datasheet for TA365670

COG2 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 20-100

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human COG2

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year

Gene Name: component of oligomeric golgi complex 2

Database Link: Entrez Gene 22796 Human

Q14746

Background: This gene encodes a subunit of the conserved oligomeric Golgi complex that is required for

maintaining normal structure and activity of the Golgi complex. The encoded protein

specifically interacts with the USO1 vesicle docking protein and may be necessary for normal

Golgi ribbon formation and trafficking of Golgi enzymes. Mutations of this gene are

associated with abnormal glycosylation within the Golgi apparatus. Alternative splicing results

in multiple transcript variants.

Synonyms: LDLC




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:




Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA365670 (COG2 Antibody) at dilution 1/25 (Original magnification: ×200)




Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA365670 (COG2 Antibody) at dilution 1/25, treated with fusion protein. (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA365670 (COG2 Antibody) at dilution 1/25 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA365670 (COG2 Antibody) at dilution 1/25, treated with fusion protein. (Original magnification: ×200)