

Product datasheet for TA365674

COMT Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB: 200-1000 WB positive control: Jurkat and Hela cell lysates IHC: 20-100 Positive control: Human thyroid cancer Predicted cell location: Cytoplasm or Nucleus
Reactivity:	Human
Host:	Rabbit
lsotype:	lgG
Clonality:	Polyclonal
Immunogen:	Fusion protein of human COMT
Formulation:	pH7.4 PBS, 0.05% NaN3, 40% Glycerol
Concentration:	lot specific
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C.
Stability:	1 year
Predicted Protein Size:	30 kDa
Gene Name:	catechol-O-methyltransferase
Database Link:	<u>Entrez Gene 1312 Human</u> <u>P21964</u>

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


GRIGENE COMT Rabbit Polyclonal Antibody – TA365674

Background:Catechol-O-methyltransferase catalyzes the transfer of a methyl group from S-
adenosylmethionine to catecholamines, including the neurotransmitters dopamine,
epinephrine, and norepinephrine. This O-methylation results in one of the major degradative
pathways of the catecholamine transmitters. In addition to its role in the metabolism of
endogenous substances, COMT is important in the metabolism of catechol drugs used in the
treatment of hypertension, asthma, and Parkinson disease. COMT is found in two forms in
tissues, a soluble form (S-COMT) and a membrane-bound form (MB-COMT). The differences
between S-COMT and MB-COMT reside within the N-termini. Several transcript variants are
formed through the use of alternative translation initiation sites and promoters.


Synonyms:

catechol-O-methyltransferase; OTTHUMP00000197749; OTTHUMP00000197750

Product images:




Gel: 8%SDS-PAGE Lysate: 40 µg Lane 1-2: Jurkat and Hela cell lysates Primary antibody: TA365674 (COMT Antibody) at dilution 1/400 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 2 minutes



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA365674 (COMT Antibody) at dilution 1/25 (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA365674 (COMT Antibody) at dilution 1/25, treated with fusion protein. (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using TA365674 (COMT Antibody) at dilution 1/25 (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using TA365674 (COMT Antibody) at dilution 1/25, treated with fusion protein. (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US