

# Product datasheet for TA365880

# **IGHA1 Rabbit Polyclonal Antibody**

# **Product data:**

Product Type:	Primary Antibodies
Applications:	IHC
Recommended Dilution:	IHC: 50-300 Positive control: Human cervical cancer Predicted cell location: Secreted, Cell membrane
Reactivity:	Human
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Immunogen:	Fusion protein of human IGHA1
Formulation:	pH7.4 PBS, 0.05% NaN3, 40% Glycerol
Concentration:	lot specific
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C.
Stability:	1 year
Gene Name:	immunoglobulin heavy constant alpha 1
Database Link:	<u>Entrez Gene 3493 Human</u> <u>P01876</u>

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

## **GRIGENE** IGHA1 Rabbit Polyclonal Antibody – TA365880


#### Background:

Constant region of immunoglobulin heavy chains. Immunoglobulins, also known as antibodies, are membrane-bound or secreted glycoproteins produced by B lymphocytes. In the recognition phase of humoral immunity, the membrane-bound immunoglobulins serve as receptors which, upon binding of a specific antigen, trigger the clonal expansion and differentiation of B lymphocytes into immunoglobulins-secreting plasma cells. Secreted immunoglobulins mediate the effector phase of humoral immunity, which results in the elimination of bound antigens (PubMed:22158414, PubMed:20176268). The antigen binding site is formed by the variable domain of one heavy chain, together with that of its associated light chain. Thus, each immunoglobulin has two antigen binding sites with remarkable affinity for a particular antigen. The variable domains are assembled by a process called V-(D)-J rearrangement and can then be subjected to somatic hypermutations which, after exposure to antigen and selection, allow affinity maturation for a particular antigen (PubMed:17576170, PubMed:20176268). Ig alpha is the major immunoglobulin class in body secretions (PubMed:2241915).

### **Product images:**



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA365880 (IGHA1 Antibody) at dilution 1/50 (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US 

Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA365880 (IGHA1 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using TA365880 (IGHA1 Antibody) at dilution 1/50 (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using TA365880 (IGHA1 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US