

## **Product datasheet for TA365941S**

## **RNF111 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: TM4 cell lysate

IHC: 50-300

Positive control: Human gastric cancer

Predicted cell location: Cytoplasm and Cell membrane

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

**Clonality:** Polyclonal

**Immunogen:** Fusion protein of human RNF111

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 109 kDa

**Gene Name:** ring finger protein 111

Database Link: Entrez Gene 54778 Human

Q6ZNA4

**Background:** The protein encoded by this gene is a nuclear RING-domain containing E3 ubiquitin ligase.

This protein interacts with the transforming growth factor (TGF) -beta/NODAL signaling pathway by promoting the ubiquitination and proteosomal degradation of negative

regulators, like SMAD proteins, and thereby enhances TGF-beta target-gene transcription. As a modulator of the nodal signaling cascade, this gene plays a critical role in the induction of mesoderm during embryonic development. Alternative splicing of this gene results in

multiple transcript variants encoding distinct isoforms.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

ARK; Arkadia; DKFZp313E0731; DKFZp686H1966; DKFZp761D081; FLJ38008

## **Product images:**



Gel: 6%SDS-PAGE Lysate: 40 µg Lane: TM4 cell lysate


Primary antibody: [TA365941] (RNF111 Antibody)

at dilution 1/200

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 1 minute



Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using [TA365941] (RNF111 Antibody) at dilution 1/50. (Original magnification: ×200)