

Product datasheet for **TA365982S**

CD48 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB: 500-2000 WB positive control: Raji, Jurkat, K562 and 231 cell, Human kidney tissue lysates IHC: 40-200 Positive control: Human esophagus cancer Predicted cell location: Cell membrane
Reactivity:	Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Fusion protein of human CD48
Formulation:	pH7.4 PBS, 0.05% NaN ₃ , 40% Glycerol
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C.
Stability:	1 year
Predicted Protein Size:	28 kDa
Gene Name:	CD48 molecule
Database Link:	Entrez Gene 962 Human P09326




[View online »](#)

Background:


This gene encodes a member of the CD2 subfamily of immunoglobulin-like receptors which includes SLAM (signaling lymphocyte activation molecules) proteins. The encoded protein is found on the surface of lymphocytes and other immune cells, dendritic cells and endothelial cells, and participates in activation and differentiation pathways in these cells. The encoded protein does not have a transmembrane domain, however, but is held at the cell surface by a GPI anchor via a C-terminal domain which maybe cleaved to yield a soluble form of the receptor. Multiple transcript variants encoding different isoforms have been found for this gene.

Synonyms:


BCM1; BLAST; BLAST1; hCD48; mCD48; MEM-102; SLAMF2; TCT.1

Product images:


Gel: 8%SDS-PAGE
 Lysate: 40 µg
 Lane 1-5: Raji
 Jurkat
 K562 and 231 cell
 Human kidney tissue lysates
 Primary antibody: [TA365982] (CD48 Antibody) at dilution 1/250
 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution
 Exposure time: 5 seconds



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using [TA365982] (CD48 Antibody) at dilution 1/40 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using [TA365982] (CD48 Antibody) at dilution 1/40, treated with fusion protein. (Original magnification: $\times 200$)