

Product datasheet for TA365995S

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

PSMA1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 1000-5000

WB positive control: Jurkat, Hela, HepG2, PC-3, Raji, 293T and SKOV3 cell lysates

IHC: 150-300

Positive control: Human esophagus cancer

Predicted cell location: Nucleus

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human PSMA1

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 30 kDa

Gene Name: proteasome subunit alpha 1

Database Link: Entrez Gene 5682 Human

P25786




Background:

The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

Synonyms:


HC2; MGC1667; MGC14542; MGC14575; MGC14751; MGC21459; MGC22853; MGC23915; NU; PROS-30; PROS30; PSC2

Product images:




Lysate: 40 µg
Lane 1-7: Jurkat
Hela
HepG2
PC-3
Raji
293T and SKOV3 cell lysates
Primary antibody: [TA365995] (PSMA1 Antibody)
at dilution 1/800
Secondary antibody: Goat anti rabbit IgG at
1/5000 dilution
Exposure time: 5 seconds

Gel: 12%SDS-PAGE



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using [TA365995] (PSMA1 Antibody) at dilution 1/180 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using [TA365995] (PSMA1 Antibody) at dilution 1/180, treated with fusion protein. (Original magnification: ×200)