

Product datasheet for TA366050

ERp72 (PDIA4) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: NIH/3T3 and RAW264.7 cell lysates

IHC: 30-150

Positive control: Human cervical cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human PDIA4

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 73 kDa

Gene Name: protein disulfide isomerase family A member 4

Database Link: Entrez Gene 9601 Human

P13667

Background: This gene encodes a member of the disulfide isomerase (PDI) family of endoplasmic

reticulum (ER) proteins that catalyze protein folding and thiol-disulfide interchange reactions.

The encoded protein has an N-terminal ER-signal sequence, three catalytically active thioredoxin (TRX) domains, two TRX-like domains and a C-terminal ER-retention sequence. This protein, when bound to cyclophilin B, enhances the rate of immunoglobulin G

intermolecular disulfide bonding and antibody assembly.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms: ERP70; ERP72

Product images:








Gel: 8%SDS-PAGE Lysate: 40 µg Lane 1-2: NIH/3T3 and RAW264.7 cell lysates Primary antibody: TA366050 (PDIA4 Antibody) at dilution 1/300 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution


Exposure time: 1 second

Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA366050 (PDIA4 Antibody) at dilution 1/50 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA366050 (PDIA4 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA366050 (PDIA4 Antibody) at dilution 1/50 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA366050 (PDIA4 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)