

Product datasheet for TA366128

H2BC15 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: TM4, K562, 293T and Jurkat cell lysates

IHC: 30-150

Positive control: Human esophagus cancer

Predicted cell location: Nucleus

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human H2BC15

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 14 kDa

Gene Name: histone cluster 1, H2bn

Database Link: Entrez Gene 8341 Human

Q99877

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H2B family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the small histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq, Aug 2015]

Synonyms:

H2B.d; H2B/d; H2BFD; HIST1H3I; MGC9388; MGC125414; MGC125415; MGC125416

Product images:



Gel: 12%SDS-PAGE Lysate: 40 µg Lane 1-4: TM4 K562 293T and Jurkat cell lysates


Primary antibody: TA366128 (H2BC15 Antibody)

at dilution 1/250

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 20 seconds



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA366128 (H2BC15 Antibody) at dilution 1/35 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA366128 (H2BC15 Antibody) at dilution 1/35, treated with fusion protein. (Original magnification: ×200)