

### **Product datasheet for TA366210**

#### Product datasneet for TA300210

## **ECH1 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: HepG2, 293T and Hela cell, Human fetal liver tissue lysates

IHC: 50-300

Positive control: Human thyroid cancer Predicted cell location: Cytoplasm

Reactivity: Human, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human ECH1

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 36 kDa

**Gene Name:** enoyl-CoA hydratase 1, peroxisomal

Database Link: Entrez Gene 1891 Human

Q13011



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




#### Background:


This gene encodes a member of the hydratase/isomerase superfamily. The gene product shows high sequence similarity to enoyl-coenzyme A (CoA) hydratases of several species, particularly within a conserved domain characteristic of these proteins. The encoded protein, which contains a C-terminal peroxisomal targeting sequence, localizes to the peroxisome. The rat ortholog, which localizes to the matrix of both the peroxisome and mitochondria, can isomerize 3-trans,5-cis-dienoyl-CoA to 2-trans,4-trans-dienoyl-CoA, indicating that it is a delta3,5-delta2,4-dienoyl-CoA isomerase. This enzyme functions in the auxiliary step of the fatty acid beta-oxidation pathway. Expression of the rat gene is induced by peroxisome proliferators.

Synonyms: HPXEL

# **Product images:**




Gel: 8%SDS-PAGE Lysate: 40 µg Lane 1-4: HepG2 293T and Hela cell Human fetal liver tissue lysates Primary antibody: TA366210 (ECH1 Antibody) at dilution 1/300 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 15 seconds



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA366210 (ECH1 Antibody) at dilution 1/50 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA366210 (ECH1 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)