

## **Product datasheet for TA366250S**

## ATP6V1C1 Rabbit Polyclonal Antibody

## **Product data:**

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Human cerebrum tissue and Human cerebella tissue lysates

IHC: 100-200

Positive control: Human tonsil Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human ATP6V1C1

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 44 kDa

Gene Name: ATPase H+ transporting V1 subunit C1

Database Link: Entrez Gene 528 Human

P21283



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that mediates acidification of intracellular compartments of eukaryotic cells. V-ATPase dependent acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A and three B subunits, two G subunits plus the C, D, E, F, and H subunits. The V1 domain contains the ATP catalytic site. The V0 domain consists of five different subunits: a, c, c', c'', and d. Additional isoforms of many of the V1 and V0 subunit proteins are encoded by multiple genes or alternatively spliced transcript variants. This gene is one of two genes that encode the V1 domain C subunit proteins and is found ubiquitously. This C subunit is analogous but not homologous to gamma subunit of F-ATPases. Previously, this gene was designated ATP6D.

Synonyms:


ATP6C; ATP6D; FLJ20057; VATC; Vma5

## **Product images:**




Gel: 8%SDS-PAGE
Lysate: 40 µg
Lane 1-2: Human cerebrum tissue and Human
cerebella tissue lysates
Primary antibody: [TA366250] (ATP6V1C1
Antibody) at dilution 1/800
Secondary antibody: Goat anti rabbit IgG at
1/5000 dilution
Exposure time: 10 seconds





Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA366250] (ATP6V1C1 Antibody) at dilution 1/100 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA366250] (ATP6V1C1 Antibody) at dilution 1/100, treated with fusion protein. (Original magnification: ×200)