

## **Product datasheet for TA366320S**

## **GSTCD Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Mouse kidney tissue lysate

IHC: 70-350

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Fusion protein of human GSTCD

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year
Predicted Protein Size: 71 kDa

**Gene Name:** glutathione S-transferase C-terminal domain containing

Database Link: Entrez Gene 79807 Human

Q8NEC7

**Background:** The function of this protein remains unknown.

**Synonyms:** DKFZp686l10174; FLJ13273




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 μg

Lane: Mouse kidney tissue lysate


Primary antibody: [TA366320] (GSTCD Antibody)

at dilution 1/1000

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 5 minutes



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA366320] (GSTCD Antibody) at dilution 1/70 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA366320] (GSTCD Antibody) at dilution 1/70, treated with fusion protein. (Original magnification: ×200)