

Product datasheet for TA366356

Histone H2A.J (H2AFJ) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Mouse brain tissue, 231, Jurkat cell lysates

IHC: 150-300

Positive control: Human esophagus cancer

Predicted cell location: Nucleus

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human H2AJ

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 14 kDa

Gene Name: H2A histone family member J

Database Link: Entrez Gene 55766 Human

Q9BTM1



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is located on chromosome 12 and encodes a replication-independent histone that is a variant H2A histone. The protein is divergent at the C-terminus compared to the consensus H2A histone family member. This gene also encodes an antimicrobial peptide with antibacterial and antifungal activity.[provided by RefSeq, Oct 2015]

Synonyms:

FLJ10903; FLJ52230; H2a/j; H2AJ; MGC921

Product images:



Gel: 12%SDS-PAGE Lysate: 40 μg

Lane 1-3: Mouse brain tissue

231

Jurkat cell lysates


Primary antibody: TA366356 (H2AJ Antibody) at

dilution 1/800

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 20 seconds



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA366356 (H2AJ Antibody) at dilution 1/160 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA366356 (H2AJ Antibody) at dilution 1/160, treated with fusion protein. (Original magnification: ×200)