

Product datasheet for TA366447S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Proteasome subunit alpha type 6 (PSMA6) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 1000-5000

WB positive control: HepG2, Hela, 293T, Jurkat and A375 cell lysates

IHC: 50-200

Positive control: Human colorectal cancer Predicted cell location: Cytoplasm and Nucleus

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human PSMA6

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 27 kDa

Gene Name: proteasome subunit alpha 6

Database Link: Entrez Gene 5687 Human

P60900






Background:

The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit. Multiple transcript variants encoding several different isoforms have been found for this gene. A pseudogene has been identified on the Y chromosome.


Synonyms:

IOTA; MGC2333; MGC22756; MGC23846; p27K; PROS-27; PROS27

Product images:




Gel: 12%SDS-PAGE
Lysate: 40 µg
Lane 1-5: HepG2
Hela
293T
Jurkat and A375 cell lysates
Primary antibody: [TA366447] (PSMA6 Antibody)
at dilution 1/900
Secondary antibody: Goat anti rabbit IgG at
1/5000 dilution
Exposure time: 90 seconds



Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using [TA366447] (PSMA6 Antibody) at dilution 1/65 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using [TA366447] (PSMA6 Antibody) at dilution 1/65, treated with fusion protein. (Original magnification: ×200)