

# **Product datasheet for TA366452**

## **Lunatic Fringe (LFNG) Rabbit Polyclonal Antibody**

### **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: K562 and HT-29 cell lysates

IHC: 50-200

Positive control: Human tonsil Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Fusion protein of human LFNG

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 42 kDa

Gene Name: LFNG O-fucosylpeptide 3-beta-N-acetylglucosaminyltransferase

Database Link: Entrez Gene 3955 Human

Q8NES3



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




#### Background:

This gene is a member of the glycosyltransferase 31 gene family. Members of this gene family, which also includes the MFNG (GeneID: 4242) and RFNG (GeneID: 5986) genes, encode evolutionarily conserved glycosyltransferases that act in the Notch signaling pathway to define boundaries during embryonic development. While their genomic structure is distinct from other glycosyltransferases, these proteins have a fucose-specific beta-1,3-N-acetylglucosaminyltransferase activity that leads to elongation of O-linked fucose residues on Notch, which alters Notch signaling. The protein encoded by this gene is predicted to be a single-pass type II Golgi membrane protein but it may also be secreted and proteolytically processed like the related proteins in mouse and Drosophila (PMID: 9187150). Mutations in this gene have been associated with autosomal recessive spondylocostal dysostosis 3.

Synonyms: SCDO3

## **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 μg

Lane 1-2: K562 and HT-29 cell lysates


Primary antibody: TA366452 (LFNG Antibody) at

dilution 1/900

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 1 minute




Immunohistochemistry of paraffin-embedded Human tonsil tissue using TA366452 (LFNG Antibody) at dilution 1/60 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human tonsil tissue using TA366452 (LFNG Antibody) at dilution 1/60, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA366452 (LFNG Antibody) at dilution 1/60 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA366452 (LFNG Antibody) at dilution 1/60, treated with fusion protein. (Original magnification: ×200)