

Product datasheet for TA366483S

PDXK Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 1000-5000

WB positive control: Mouse brain tissue, MCF-7 cell lysates

IHC: 50-200

Positive control: Human prostate cancer Predicted cell location: Cytoplasm and Nucleus

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein of human PDXK

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 35 kDa

Gene Name: pyridoxal (pyridoxine, vitamin B6) kinase

Database Link: Entrez Gene 8566 Human

O00764

Background: The protein encoded by this gene phosphorylates vitamin B6, a step required for the

conversion of vitamin B6 to pyridoxal-5-phosphate, an important cofactor in intermediary metabolism. The encoded protein is cytoplasmic and probably acts as a homodimer.

Alternatively spliced transcript variants have been described, but their biological validity has

not been determined.

Synonyms: C21orf97; C21orf124; DKFZp566A071; FLJ21324; FLJ31940; FLJ37311; MGC15873; MGC31754;

MGC52346; PKH; PNK; PRED79




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:




Gel: 8%SDS-PAGE
Lysate: 40 µg
Lane 1-2: Mouse brain tissue
MCF-7 cell lysates
Primary antibody: [TA366483] (PDXK Antibody) at
dilution 1/2000
Secondary antibody: Goat anti rabbit IgG at
1/5000 dilution
Exposure time: 20 seconds



Immunohistochemistry of paraffin-embedded Human prostate cancer tissue using [TA366483] (PDXK Antibody) at dilution 1/75 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human prostate cancer tissue using [TA366483] (PDXK Antibody) at dilution 1/75, treated with fusion protein. (Original magnification: ×200)