

# Product datasheet for TA366510S

**Primary Antibodies** 

WB positive control: 293T and K562 cell lysates

Positive control: Human prostate cancer

Predicted cell location: Nucleus

Fusion protein of human SUV39H2

Antigen affinity purification

Entrez Gene 79723 Human

pH7.4 PBS, 0.05% NaN3, 40% Glycerol

suppressor of variegation 3-9 homolog 2

IHC, WB

IHC: 50-200

Human, Mouse

Rabbit

Polyclonal

Unconjugated Store at -20°C.

1 year

47 kDa

Q9H5I1

lgG

## SUV39H2 Rabbit Polyclonal Antibody

Recommended Dilution: WB: 500-2000

### **Product data:**

**Product Type:** 

**Applications:** 

**Reactivity:** 

Host:

Isotype:

**Clonality:** 

Immunogen:

Formulation:

**Purification:** 

**Conjugation:** 

Gene Name:

Database Link:

**Predicted Protein Size:** 

Storage: Stability: OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


#### **GRIGENE** SUV39H2 Rabbit Polyclonal Antibody – TA366510S

Background: Histone methyltransferase that specifically trimethylates 'Lys-9' of histone H3 using monomethylated H3 'Lys-9' as substrate. H3 'Lys-9' trimethylation represents a specific tag for epigenetic transcriptional repression by recruiting HP1 (CBX1, CBX3 and/or CBX5) proteins to methylated histones. Mainly functions in heterochromatin regions, thereby playing a central role in the establishment of constitutive heterochromatin at pericentric and telomere regions. H3 'Lys-9' trimethylation is also required to direct DNA methylation at pericentric repeats. SUV39H1 is targeted to histone H3 via its interaction with RB1 and is involved in many processes, such as cell cycle regulation, transcriptional repression and regulation of telomere length. May participate in regulation of higher-order chromatin organization during spermatogenesis. Recruited by the large PER complex to the E-box elements of the circadian target genes such as PER2 itself or PER1, contributes to the conversion of local chromatin to a heterochromatin-like repressive state through H3 'Lys-9' trimethylation.

Synonyms:


FLJ23414; KMT1B

#### **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 μg Lane 1-2: 293T and K562 cell lysates Primary antibody: [TA366510] (SUV39H2 Antibody) at dilution 1/700 Secondary antibody: Goat anti rabbit IgG at 1/5000 dilution Exposure time: 3 minutes

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemistry of paraffin-embedded Human prostate cancer tissue using [TA366510] (SUV39H2 Antibody) at dilution 1/70 (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human prostate cancer tissue using [TA366510] (SUV39H2 Antibody) at dilution 1/70, treated with fusion protein. (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US