

Product datasheet for TA366560S

HEXA Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies Applications: IHC, WB Recommended Dilution: WB: 500-2000 WB positive control: Jurkat, PC3 and HT-29 cell lysates IHC: 100-200 Positive control: Human tonsil Predicted cell location: Cytoplasm **Reactivity:** Human, Mouse, Rat Host: Rabbit Isotype: lgG **Clonality:** Polyclonal Immunogen: Fusion protein of human HEXA Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol **Purification:** Antigen affinity purification **Conjugation:** Unconjugated Store at -20°C. Storage: Stability: 1 year **Predicted Protein Size:** 61 kDa Gene Name: hexosaminidase subunit alpha Database Link: Entrez Gene 3073 Human

<u>P06865</u>

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

MEXA Rabbit Polyclonal Antibody – TA366560S


Background:This gene encodes a member of the glycosyl hydrolase 20 family of proteins. The encoded
preproprotein is proteolytically processed to generate the alpha subunit of the lysosomal
enzyme beta-hexosaminidase. This enzyme, together with the cofactor GM2 activator protein,
catalyzes the degradation of the ganglioside GM2, and other molecules containing terminal
N-acetyl hexosamines. Mutations in this gene lead to an accumulation of GM2 ganglioside in
neurons, the underlying cause of neurodegenerative disorders termed the GM2
gangliosidoses, including Tay-Sachs disease (GM2-gangliosidosis type I). Alternative splicing
results in multiple transcript variants, at least one of which encodes a preproprotein that is
proteolytically processed.

Synonyms: beta-N-acetylhexosaminidase; MGC99608; N-acetyl-beta-glucosaminidase; TSD

Product images:




Gel: 8%SDS-PAGE Lysate: 40 µg Lane 1-3: Jurkat PC3 and HT-29 cell lysates Primary antibody: [TA366560] (HEXA Antibody) at dilution 1/700 Secondary antibody: Goat anti rabbit IgG at 1/5000 dilution Exposure time: 1 minute



Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA366560] (HEXA Antibody) at dilution 1/100 (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA366560] (HEXA Antibody) at dilution 1/100, treated with fusion protein. (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA366560] (HEXA Antibody) at dilution 1/100 (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA366560] (HEXA Antibody) at dilution 1/100, treated with fusion protein. (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US