

### **Product datasheet for TA366822**

## **LAYN Rabbit Polyclonal Antibody**

### **Product data:**

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Jurkat and K562 cell lysates

IHC: 50-100

Positive control: Human esophagus cancer Predicted cell location: Cell membrane

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human LAYN

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year
Predicted Protein Size: 43 kDa
Gene Name: layilin

Database Link: Entrez Gene 143903 Human

Q6UX15

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn






#### Background:

Layilin is a recently cloned ~55-kDa membrane-binding partner for talin (Borowsky and Hynes, 1998 ) that is widely expressed in different cell types and tissue extracts. It is found in peripheral ruffles of spreading cells and is recruited to membrane ruffles in cells induced to migrate in in vitro wounding experiments. Layilin colocalizes with talin in ruffles and binds to talin's ~50-kDa head domain (amino acids 280–435). Other parts of talin (the ~220-kDa tail fragment) can bind  $\beta$ -integrin cytoplasmic tails, vinculin, and F-actin and hence form an integrin-cytoskeleton linkage at sites of cell-substratum contact. Layilin's extracellular domain is homologous with the carbohydrate-recognition domains (CRD) of C-type lectins. Receptor for hyaluronate.

**Synonyms:** FLJ30977; FLJ31092; layilin

# **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 μg

Lane 1-2: Jurkat and K562 cell lysates


Primary antibody: TA366822 (LAYN Antibody) at

dilution 1/700

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 2 minutes



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA366822 (LAYN Antibody) at dilution 1/50 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TA366822 (LAYN Antibody) at dilution 1/50, treated with synthetic peptide. (Original magnification: ×200)