

## **Product datasheet for TA366966S**

## **SMAD1 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 25-100

Positive control: Human thyroid cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen:Synthetic peptide of human SMAD1Formulation:pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year

**Gene Name:** SMAD family member 1

**Database Link:** Entrez Gene 4086 Human

Q15797

**Background:** The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the

gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. SMAD proteins are signal transducers and transcriptional modulators that

mediate multiple signaling pathways. This protein mediates the signals of the bone

morphogenetic proteins (BMPs), which are involved in a range of biological activities including cell growth, apoptosis, morphogenesis, development and immune responses. In response to BMP ligands, this protein can be phosphorylated and activated by the BMP receptor kinase. The phosphorylated form of this protein forms a complex with SMAD4, which is important for

its function in the transcription regulation.

Synonyms: BSP-1; BSP1; CMT2D; GARS; HMN5; hSMAD1; IV4-1; IV41; MADH1; MADR1




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA366966] (SMAD1 Antibody) at dilution 1/40 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA366966] (SMAD1 Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: ×200)